

A Survey for Low-mass Stars and Brown Dwarfs in the Upper-Scorpius OB Association

David R. Ardila¹, Eduardo L. Martín², Gibor Basri¹

Received _____; accepted _____

DRAFT - To appear in AJ

¹Astronomy Dept., Univ. of California, Berkeley, CA 94720,
e-mail: ardila@garavito.berkeley.edu, basri@soleil.berkeley.edu

²Division of Geological and Planetary Sciences, Caltech 150-21, Pasadena, CA 91125,
e-mail: ege@gps.caltech.edu

ABSTRACT

The Upper-Scorpius association is the OB association nearest to the Sun (145 pc). Its young age (~ 5 Myr) makes it an ideal place to search for low-mass stars and brown dwarfs, as these objects should be relatively bright. We have performed a photometric search for the low-mass members of the association, using the R, I, and Z filters. The completeness limit is $I \sim 18.5$ and the saturation limit is $I \sim 13$. We obtain 138 candidate members, covering nearly the entire M spectral type range. We find an excess of brown dwarf candidates over the number predicted by a Miller-Scalo Initial Mass Function. In addition, we have performed infrared imaging and low resolution optical spectroscopy of selected candidates. We find that the infrared observations confirm the spectral types obtained with the optical photometry. Furthermore, we find $H\alpha$ in emission in 20 of the 22 objects observed spectroscopically. As $H\alpha$ is an indicator of youth, we believe that these 20 objects may belong to the association. One of them, UScoCTIO 128 has a very strong and constant $H\alpha$ line (equivalent width: ~ 130 Å), and its position in the color magnitude diagram suggests that it is a brown dwarf with mass equal to $0.02 M_{\odot}$. Confirmation of this and the other candidates will have to wait for higher resolution observations that can reveal spectroscopic mass indicators like Li I and gravity indicators, such as K I and the subordinate lines of Na I.

Subject headings: stars: low-mass; stars: brown dwarfs; clusters: Upper-Scorpius

1. Introduction

OB associations and gravitationally unbound clusters are likely to be the dominant birthplaces for the low-mass field star population (Preibisch & Zinnecker 1999). Furthermore, they provide an opportunity to study very young low-mass stars and brown dwarfs, since these objects should be relatively bright in very young regions. Besides their intrinsic interest, the low-mass population of OB associations can provide constraints on the shape of the Initial Mass Function (IMF) at low masses. There are indications (Bouvier et al. 1998) that the shape of this IMF may deviate from the simple Miller-Scalo (Miller & Scalo 1979) at masses less than $0.1 M_{\odot}$.

While the low-mass stellar content of T associations (such as Taurus-Auriga, see Briceño et al. 1998 and Kenyon & Hartmann 1995) or very young clusters (such as IC 348, see Herbig 1998) seems well known, not much is known about the low-mass stellar content of OB associations. Part of the difficulty is due to the fact that most of the low-mass pre-main sequence (PMS) stars cannot be easily distinguished from normal field stars in the huge area on the sky (several hundred square degrees) covered by nearby OB associations. Only Classical T-Tauri stars can be found easily by their strong $H\alpha$ emission, using objective-prism surveys. X-ray observations have also proved to be an efficient way of distinguishing PMS stars from older field stars (see below).

The Scorpius-Centaurus association is the OB association nearest to the Sun. It contains several hundred B stars arranged in three subgroups: Upper-Scorpius, Upper Centaurus-Lupus and Lower Centaurus-Crux. Upper-Sco ($l \simeq 354^{\circ}$, $b \simeq 20^{\circ}$) is the youngest subgroup. Its lack of dense molecular material and deeply embedded young stellar objects indicates that the process of star formation has ended. The area is free of dense gas and clouds, and the association members show only moderate extinction ($A_V < 2$ mag).

Hipparcos data has been used to identify 120 association members, including 49 B

stars and 34 A stars. According to these data, Upper-Sco is 145 pc away from the Sun and has a size of $\sim 130\text{deg}^2$ (De Zeeuw et al. 1999). Age determinations based on the upper (de Zeeuw & Brand 1985; de Geus et al. 1989) and lower (Preibisch & Zinnecker 1999) mass range derive a value of 5 Myr. However, there have been age determinations that seem to suggest that many of the low-mass cluster members could be 10 Myr old (Martín 1998; Frink 1999).

The first search for low-mass members of Upper-Sco was performed by Walter et al. (1994). They obtained photometry and spectroscopy for the optical counterparts of X-ray sources detected in seven Einstein fields and classified 28 objects as low-mass PMS stars. Two large-scale surveys have been performed recently. The first was conducted by Kunkel (1999) who observed optical counterparts of more than 200 *ROSAT* All-Sky Survey (RASS) X-ray sources in a $\sim 60\text{deg}^2$ area in Upper-Scorpius and Upper Centaurus-Lupus. The other study was a spectroscopic survey for PMS stars in a 160deg^2 area by Preibisch et al. (1998). A number of further searches have been performed, all focused on small subregions within the association (Meyer et al. 1993; Sciortino et al. 1998; Martín 1998). A study of the history of star formation in Upper-Sco has been published by Preibisch & Zinnecker (1999). The authors obtained R and I photometry of association members to about $I \sim 12.8$ and $R \sim 14$.

With the goal of extending the low-mass sequence of the association, we have obtained photometry for the association in the R, I, and Z filters. We have also observed selected member candidates in the J and H filters and spectroscopically. Our search starts at $I \sim 13$ and it is therefore complementary to Preibisch & Zinnecker (1999). Section 2 describes the observations and section 3 outlines the results.

2. Observations

A summary of all observations is in Table 1.

We obtained photometry for eight fields of 80 by 80 arcmins in Upper-Sco using the 60cm Michigan Curtis-Schmidt telescope at CTIO. We therefore cover $\sim 10\%$ of the association. Figure 1 shows the location of the fields. We observed the fields in the R, I, and Z filters. Raw frames were reduced within the IRAF¹ environment, using the CCDRED package. The images were bias-subtracted and flat-fielded. The photometry was obtained using the PSF fitting routines from IRAF. As stars are undersampled (the average FWHM of a star is ~ 1.5 pixels) the errors in the photometry are dominated by centering errors. We obtained magnitudes of more than 180,000 stars.

The completeness limits for each filter are: $R \sim 19$ magnitudes, $I \sim 18.5$ magnitudes, $Z \sim 18.5$ magnitudes. All our fields saturate at ~ 13 mag. in all the filters. Figures 2ab show the color-magnitude diagrams for R-I and I-Z. Also shown are the completeness limits and saturation limits. We select as preliminary candidates those objects that lie to the right of the Leggett (1992) main sequence in both color-magnitude diagrams and below the saturation limit in the I vs. I-Z color-magnitude diagram. This assumption will most certainly increase the number of candidates that are not cluster members. The best way of doing the selection would be to include the errors due to photometry and undetected binaries in the estimation of the number of objects in a band around the 5 Myr isochrones. However, given that we do not have *a priori* information about reddening, it seems safer to begin by assuming that every object to the right of the Leggett (1992) sequence will be a

¹IRAF is distributed by National Optical Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc., under contract with the National Science Foundation

member of the cluster.

With this selection method we also take account of the saturation of the R-I color for low-mass stars. As has been shown by Bessell (1991), the R-I color saturates at $R - I \sim 2.4$ (due to molecular absorption in the stellar photosphere) and becomes bluer for cooler objects. Therefore, a selection based only on a band around the 5 Myr isochrones would miss the very low-mass objects. Figure 3 shows the I vs. R-I color-magnitude diagrams for the resulting candidates. As can be seen from Figure 3, the saturation limit in the Z filter affects the selection of candidates brighter than $I \sim 13.5$ which corresponds to $\sim M2$. In the lower limit, we will miss objects with $R - I > 2.2$.

We obtain 138 candidates, listed in Table 2 (finding charts can be obtained by contacting the authors). If all candidates belonged to the cluster then we would neatly cover all the range of M stars. As mentioned in the introduction, Preibisch & Zinnecker (1999) obtained R and I photometry of association members to about $I \sim 12.8$ and $R \sim 14$. Our search therefore begins where theirs ended.

We have complemented the optical observations with infrared J and H observations of selected candidates. To perform these observations we used the Cerro Tololo Infrared Imager (CIRIM) at CTIO. Raw frames were reduced within the IRAF environment, following the procedure outlined by Joyce (1992). We were able to observe only 9 objects, due to bad weather and instrumental problems. Table 3 details the results.

Using the red arm of the KAST spectrograph at the Lick 3m telescope (which covers the range from 5000 to 10000 Å), we observed selected bright candidates in low resolution (Grating 300/7500, which gives $\Delta\lambda \sim 11$ Å of resolution). Raw images were reduced within the IRAF environment, using standard tools to perform flat-fielding, optimal extraction, wavelength calibration and response correction of the spectra. The spectra were not corrected for telluric absorption.

Because of the low declination of the cluster, observations from Lick Observatory must be made through a large airmass. This, together with the low resolution of the observations and the variable fringing of the spectrograph in the red arm, make it very difficult to comment on gravity sensitive lines that lie on regions of telluric absorption, such as the K I resonance doublet ($\lambda\lambda 7665, 7699$) or the subordinate lines of the Na I doublet ($\lambda\lambda 8183, 8195$). However, the spectral resolution is good enough to identify whether or not a star has $H\alpha$ in emission. We have used the I3 index defined by Martín & Kun (1996) and the VO index defined by Kirkpatrick, Henry & Simons (1995) to find the spectral type of the Lick stars. These indexes have the advantage of being based on flux ratios that are close to each other. Therefore, they are not very sensitive to reddening. The spectral type obtained from the spectroscopic indexes can be compared with the spectral type from the colors to obtain the reddening. The results are shown in Table 3. The spectral types derived from the spectroscopy confirm the spectral types from the colors.

3. Discussion

3.1. Contamination by other sources

Photometric observations as a way to select cluster candidates are susceptible to contamination by foreground and background objects. There are four kinds of objects that may appear above the main sequence in the color-magnitude diagram: background giants, background galaxies, reddened background stars, and foreground low-mass stars.

From the results of Kirkpatrick et al. (1994) it is possible to show that the number of contaminating giants is negligible ($< 5\%$ for all spectral types). Contamination by galaxies is not important for the region of interest in the color magnitude diagram (Bouwens et al. 1998ab).

To estimate the contamination due to reddened background stars we use the maximum reddening towards the Upper-Sco region, which is $A_V \sim 2$ (Schlegel, Finkbeiner & Davis, 1998). Using the density of background stars (from the observed population of background stars under the Leggett sequence in the color-magnitude diagrams) we find that at most 25 candidates may be background stars: 7 before M4, 10 between M4 and M5, 8 after M5. These will preferentially lie close to the Leggett sequence.

Another source of contamination is foreground field M-stars in the cluster line of sight. Using the luminosity function derived by Kirkpatrick et al. (1994) we find that there should be 10 field stars between M4 and M5, 3 between M5 and M6, 6 between M6 and M7, and 15 between M7 and M9. Before M4 we expect a lower limit of 15 field stars.

3.2. Spectroscopic observations

Low resolution spectroscopic observations of 22 candidates were made with the idea of determining cluster membership. As was mentioned before, the observations are not detailed enough to give information about gravity, but they can provide information about activity in the form of the $H\alpha$ line. As has been shown by Prosser, Stauffer & Kraft (1991), activity decreases with age, and therefore a strong $H\alpha$ line in emission is an indicator of youth. Prosser et al. (1991) and Liebert et al. (1992) have also shown that activity increases with spectral type, starting about M1 for field dwarfs. The $H\alpha$ equivalent width reaches $-12 \text{ \AA}$ at about M9.

We find $H\alpha$ in 20 of the 22 objects observed spectroscopically. The values of the equivalent width are in Table 3. In Table 3 we also compare the spectral types determined from the spectra to those determined from CIRIM observations (I-J) when possible. As the Table shows, the results between the measurements are consistent, which confirms

the accuracy of the photometry. Figure 4 shows the traces of five representative spectra, uncorrected for telluric absorption but corrected for reddening.

Figure 5 shows the comparison of the measured $H\alpha$ equivalent widths in Upper-Sco with those in the σ -Orionis (3-7 Myr, Béjar et al. 1999) and the α -Persei (60 - 90 Myr, Stauffer et al. 1999, Basri & Martín 1999) clusters. Also shown is the envelope of $H\alpha$ equivalent widths for field stars (Prosser et al. 1991; Liebert et al. 1992). Overall, the Upper-Sco values lie above those of α -Per and below those of σ -Ori. 9 of the Upper-Sco objects are below the envelope defined by field stars, which means that their $H\alpha$ strength is consistent with them not being cluster members. However, the smoothness of the envelope is misleading. The original data for field stars shows that the maximum $H\alpha$ equivalent width for each spectral type has a fair amount of scattering. Furthermore, spectroscopic observations of the young σ -Orionis cluster have shown that the three objects below the field-star envelope are likely to be cluster members. These arguments show that the presence of $H\alpha$ alone is not enough to confirm or deny the status of a candidate as a cluster member.

Figure 6 shows the color-magnitude diagram for the objects observed spectroscopically. All the colors and magnitudes have been corrected for reddening, if known from the spectroscopy. The errors in each axis are $\sim \pm 0.1$ magnitudes. We have included in Figure 6 the isochrones calculated by D’Antona & Mazzitelli (1994). The translation from their Luminosity- T_{eff} calculations to I and R-I involves a color- T_{eff} scale and bolometric corrections. We have used the transformations given by Bessell, Castelli & Plez (1998) and the colors for low-mass stars from Kirkpatrick & McCarthy (1994). As Bessell et al. (1998) give bolometric corrections only for dwarf stars and the Kirkpatrick & McCarthy (1994) colors are from field stars, the color-magnitude isochrones in Figure 6 suffer from considerable uncertainties. This is another reason to base the selection of candidates in terms of the Leggett (1992) main sequence.

Besides D’Antona & Mazzitelli (1994), Burrows et al. (1997) and Baraffe et al. (1998) have published low-mass isochrones. Burrows et al. (1997) provide luminosities and effective temperatures, and so the translation to observable quantities suffer from the same problems as indicated above. The predicted masses for each color are smaller by about 50%, compared to those of D’Antona & Mazzitelli (1994). Baraffe et al. (1998) provide isochrones in the color-magnitude space but they are too blue, touching the observed Leggett main sequence already at 5 Myr. A comparison of the models of the three groups can be found in Béjar, Zapatero-Osorio & Rebolo (1999). This comparison shows that there may still be systematic errors in the R,I isochrones.

Even taken into account the measurement errors, the objects are scattered around various isochrones. Assuming they all belong to the cluster, it is not clear from these observations what would be the correct age of the association, because even those objects with strong $H\alpha$ (e.g. objects above the $H\alpha$ envelope for the field) do not all fall on a single isochrone. If one believes the 5 Myrs estimate for the age of the cluster (Preibisch & Zinnecker 1998), the scatter has to be explained by other means. If some of the spectroscopic objects were unresolved binaries, their magnitudes would have to be increased. This would work for the latest types of objects, but not for earliest types. As mentioned above, errors in the theoretical isochrones are also a possibility. However, it does not seem likely that any realistic adjustment in the models will make all the data points lie on the same isochrone. All these arguments point to the conclusion that the scatter is probably caused by more than one factor. Without follow-up spectroscopy (see below) it is not possible to make a stronger statement.

Of the 11 objects with a spectral type between M4 and M5 we find $H\alpha$ in all except two. We expect 40% contamination in this bin, and given the small-number statistics involved, our findings of two contaminating stars are consistent with this estimate, assuming

that all the stars with $H\alpha$ in emission are indeed cluster members. We found $H\alpha$ on the eight stars observed with spectral types between M5 and M6. The expected contamination is about 10%. The fact that we did not find any clear non-cluster member is consistent with the estimate. In other words, assuming that the objects with $H\alpha$ in emission are cluster members is consistent with the contamination estimates.

3.3. The Initial Mass Function

From the D’Antona & Mazitelli (1994) models we find that the substellar limit for this association should be at $I \sim 14.6$, $R - I \sim 2.1$, $\sim M6$. This is not an accurate number, given the uncertainties in the color- T_{eff} scale and the bolometric corrections mentioned before. However, if one accepts this estimate, we should have 10 brown dwarfs in our spectroscopic sample, assuming that all the $H\alpha$ emitters belong to the association.

One can use the Miller-Scalo IMF (Miller & Scalo 1979) centered in $0.1 M_{\odot}$ to estimate the expected number of low-mass stars and brown dwarfs in our sample. Using the 34 A stars found by Hipparcos (De Zeeuw et al. 1999) we obtain that there should be 650 stars with masses between 0.07 and $0.5 M_{\odot}$, and about 600 brown dwarfs with masses between 0.005 and $0.07 M_{\odot}$. We are only complete to $0.03 M_{\odot}$: there should be 30 brown dwarfs between 0.03 and $0.07 M_{\odot}$. Given that we are covering 10% of the cluster, we expect 65 M stars and 60 brown dwarfs, with 3 brown dwarfs having masses between 0.03 and $0.07 M_{\odot}$. In the whole sample, we find ~ 20 objects with masses between 0.03 and $0.07 M_{\odot}$ (the number of expected contaminants is ~ 10) and ~ 90 objects with masses greater than $0.03 M_{\odot}$ (the number of expected contaminants is ~ 20). The number of M Stars (70) is similar to that predicted by the Miller-Scalo IMF. The number of brown dwarf candidates that we are finding here is a little high (3 predicted compared to 10 after taking account of the contamination). As the recent work by Paresce & De Marchi (1999) shows, most

estimations of the IMF assume a log-normal distribution, generally centered in masses larger than $0.1 M_{\odot}$. Using one of such distributions would decrease even more the number of predicted brown dwarfs in our sample.

This discrepancy between observed and predicted number of brown dwarfs may be due to an underestimation of the number of contaminants. The study by Kirkpatrick et al. (1994) mentioned in the section about contamination measures stellar populations towards the galactic poles. Such populations may not be representative of the line of sight towards Upper-Scorpius. The study by Guillout et al. (1998) finds a population of young late-type stars (down to M2) in nearby Lupus distributed along the so-called ‘Gould Belt’. It is possible that the Belt population is contaminating our sample as it crosses near Upper-Scorpius, even though it is difficult to understand why it affects only the brown dwarfs range and not the low-mass stars range. Alternatively, the discrepancy between observed and expected number of brown dwarfs may reflect a real increase in the IMF at low masses, similar to that described by Bouvier et al. (1998). It is not possible to make a stronger statement about the IMF without follow-up spectroscopy of all the candidates.

3.4. Lithium Burning

As models show, the transition between lithium destruction and preservation in stellar photospheres occurs at higher masses for younger clusters. At the age of Upper-Sco (5 Myr) neither very low-mass stars nor brown dwarfs have had sufficient time to reach the core temperatures necessary to start lithium burning (D’Antona & Mazzitelli 1997; Soderblom et al. 1998). We therefore plan to search for lithium in these candidates, as they should have it if they are cluster members. As the cluster is very young, the so-called ‘Lithium Test’ (Magazzu, Martín & Rebolo 1993), the use of the substellar lithium boundary to date the cluster, cannot be applied in Upper-Sco. However, the amount of lithium in the

photosphere of early M stars could be used to date the association, as an M0 member of a 10 Myr cluster should have about 4 times less lithium than if the cluster were 5 Myr old (Soderblom et al. 1998). Also, as always, an object with spectral type later than M7 and showing lithium in the photosphere should be a brown dwarf (Basri 1998). Therefore, even though there is no lithium boundary in the cluster, lithium is still an important age and mass diagnostic.

Recently, Béjar et al. (1999) have suggested that the deuterium boundary could be used as another way to date the cluster. For the Upper-Sco association we expect the deuterium boundary to be located at $0.04M_{\odot}$ and therefore some of our reddest candidates could still have deuterium in their atmospheres. However, detecting of deuterium abundances poses considerable challenges from the observational point of view.

UScoCTIO 128 is a very interesting object: it shows strong $H\alpha$ emission, and its position in the color magnitude diagram indicates a mass of $0.02 M_{\odot}$. For comparison, the models from Burrows et al. (1997) give a mass even smaller, of about $0.015 M_{\odot}$. We have two measurements of the $H\alpha$ equivalent width (separated by a month and taken with the same instrument in the same configuration), and the value is the same within the errors. This indicates that the strong equivalent width is probably not a flare. In this respect UScoCTIO 128 is very similar to S Ori 45, found by Béjar et al. (1999). More spectroscopic observations are needed before we can confirm UScoCTIO 128 as the lowest mass member yet of the Upper-Sco OB association.

4. Conclusions

We have conducted a photometric search for the low-mass members of the Upper-Scorpius OB association, using the R, I, and Z filters. Completeness limits are $R \sim 19$,

$I \sim 18.5$, $Z \sim 18.5$. Our search covers $\sim 10\%$ of the association. The photometry comfortably crosses the substellar limit for the cluster, situated at $I \sim 14.6$, $R - I \sim 2.1$, $\sim M6$. This is the first survey to sample the low-mass region of Upper-Sco. We find 138 candidate members of the cluster. Contamination by non-cluster members (mainly foreground M stars and background reddened stars) has been estimated to be 59 objects.

Follow-up observations using infrared images and low resolution spectroscopy confirm the optical photometry of a reduced sample of the candidates. Of 22 objects observed spectroscopically, 20 have $H\alpha$ in emission, an indicator of young age. Comparisons between the $H\alpha$ equivalent widths found in other clusters and those in Upper-Sco indicate that 11 of those 20 objects may be members of the association, as those 11 object have stronger $H\alpha$ than expected for low-mass field stars. However, it is possible that all the 20 objects are indeed association members. The objects with strong $H\alpha$ do not all fall on a single isochrone. This may be due to the presence of unresolved binaries, to contamination from field $H\alpha$ emitters or to an intrinsic age spread in the cluster.

Using the Miller-Scalo IMF centered on $0.1 M_{\odot}$ we estimate the number of objects with masses between 0.07 and $0.3 M_{\odot}$ as 3 times less than what is observed. This may be due to an underestimation on the number of contaminants affecting this mass bin, due for example to a contribution of low-mass objects by the Gould Belt. On the other hand, similar excesses in the low-mass populations have been observed in other young clusters, like the Pleiades, and may point to a departure of the IMF from a simple log-normal form. A stronger statement about the IMF will have to wait until we have more cluster membership diagnostics.

The spectroscopic observations do not have high enough resolution to observe gravity sensitive lines, such as the resonant transitions of K I or the subordinate lines of Na I. Therefore, they can not be used to distinguish between young low-mass objects and

main-sequence objects.

We find a very interesting object, UScoCTIO 128 ($\sim$ M7) with very strong H α (equivalent width of ~ -130 Å) and an estimated mass of $0.02 M_{\odot}$. If this object is a member of the cluster, it would be one of the lowest-mass brown dwarfs known to date.

Full confirmation of the membership of these objects will have to wait for higher resolution spectroscopy that can observe Li I, K I, and Na I. As has been suggested by other groups, these very young objects provide a unique opportunity to study depletion of very light elements such as deuterium. Perhaps in the future studies of deuterium depletion will take the place of lithium depletion as a precise method of determining the ages of very young clusters.

We would like to thank Victor Béjar for invaluable help in collecting the CTIO observations and for many stimulating discussions concerning brown dwarfs. Thanks are also due to Debi Howell-Ardila, who edited the manuscript for language. We acknowledge the support of the National Science Foundation through grant number AST96-18439. EM acknowledges support from the NASA Origins program.

REFERENCES

- Baraffe, I., Chabrier, G., Allard, F., & Hauschildt, P. H. 1998, *A&A*, 337, 403
- Basri, G. 1998, in ASP Conf. Ser. 134, *Brown Dwarfs and Extrasolar Planets*, ed. R. Rebolo, E. L. Martín & M. R. Zapatero-Osorio (San Francisco: ASP), 394
- Basri, G. & Martín, E. L. 1999, *ApJ*, 510, 266
- Béjar, V. J. S., Zapatero-Osorio, M. R. & Rebolo, R. 1999, *ApJ*, 521, 671
- Bessell, M. S., 1991, *AJ*, 101, 662
- Bessell, M. S., Castelli, F. & Plez, B. 1998, *A&A*, 333, 231
- Bouvier J. et al. 1998, 336, 490
- Bouwens, R.J., Broadhurst, T.J. & Silk, J. 1998a, *ApJ*, 506, 557
- Bouwens, R.J., Broadhurst, T.J. & Silk, J. 1998b, *ApJ*, 506, 579
- Briceño, C., Hartmann, L., Stauffer, J. & Martín, E. L. 1999, 115, 2074
- D’Antona, F. & Mazzitelli, I. 1994, *ApJS*, 90, 467
- D’Antona, F. & Mazzitelli, I. 1997, ‘Evolution of low-mass stars’, in “Cool stars in Clusters and Associations”, eds. G. Micela and R. Pallavicini, *Mem.S.A.It.*, 68, 807
- de Geus, E. J., De Zeeuw, P. T. & Lub, J. 1989, *A&A*, 216, 44
- De Zeeuw, P. T. & Brand, J. 1985, in *Birth and Evolution of Massive Stars and Stellar Groups*, ed. H. van den Woerden & W. Boland (Dordrecht: Reidel), 95
- De Zeeuw, P. T., Hoogerwerf, R., De Bruijne, J. H. J., Brown, A. G. A. & Blaauw, A. 1999, *AJ*, 117, 354

- Frink, S. 1999, PhD Thesis, Ruprecht-Karls-Universitt Heidelberg
- Guillout, P., Sterzik, M. F., Schmitt, J. H. M. M., Motch, C. & Neuhausser, R. 1998, A&A, 337,113
- Herbig, G. 1998, ApJ, 497, 736
- Joyce, R. 1992, in ASP Conf. Ser. 23, Astronomical CCD observing and reduction techniques, ed. S. B. Howell (San Francisco: ASP), 258
- Kenyon, S. J. & Hartmann, L. W. 1995, ApJS, 101,117
- Kirkpatrick, J. D., McCarthy, D. W. 1994a, AJ, 107, 333
- Kirkpatrick, J. D., McGraw, J. T., Hess, T. R., Liebert, J. & McCarthy, D. W. 1994, ApJS, 94, 749
- Kirkpatrick, J. D., Henry, T. J. & Simons, D. A. 1995, 109, 797
- Kunkel, M. 1999, PhD Thesis, Universität Würzburg
- Leggett, S. K. 1992, ApJS, 82, 351
- Liebert, J., Saffer, R. A., Norsworthy, J., Giampapa, M. S. & Stauffer, J. R. 1992, in ASP Conf. Ser. 26, Seventh Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun, ed. M. S. Giampapa & J. A. Bookbinder (San Francisco: ASP), 282
- Magazzu, A., Martín, E. L. & Rebolo, R. 1993, ApJ, 404, 17
- Martín, E. L. & Kun, M. 1996, A&AS, 116, 467
- Martín, E. L. 1998, AJ, 115, 351
- Meyer, M. R., Wilking, B. A. & Zinnecker, H. 1993, AJ, 105, 619

Miller, G. & Scalo, J. 1979, ApJS, 41 513

Paresce, F. & De Marchi, G. 1999, ApJ, in press

Preibisch, Th., Guenther, E., Zinnecker, H., Sterzik, M., Frink, S. & Röser, S. 1998, A&A, 333, 619

Preibisch, Th. & Zinnecker, H. 1999, AJ, 117, 238

Prosser, C. F., Stauffer, J., Kraft, R. P. 1991, AJ, 101, 136

Schlegel, D. J., Finkbeiner, D. P. & Davis, M. 1998, ApJ, 500, 525

Sciortino, S., Damiani, F., Favata, F. & Micela, G. 1998, A&A, 332, 825

Soderblom, D. R. et al. 1998, ApJ, 498, 385

Stauffer, J. et al. 1999, ApJ, in press

Walter, F. M., Vrba, F. J., Mathieu, R. D., Brown, A. & Myers, P. C. 1994, AJ, 107, 692

Figure 1: Location of the observed fields in Upper-Scorpius. Each field is 80' by 80'. The black dots indicate Hipparcos members (De Zeeuw et al. 1999).

Figure 2: (a) I vs. R-I color-magnitude diagram for all the objects observed. The results of the photometry are represented by small dots. The lower dashed line is the completeness limit. The solid line is the Leggett (1992) main sequence. (b) Same diagram for I vs. I-Z. The upper dashed line is the saturation limit.

Figure 3: Color-magnitude diagram for the selected candidates. The solid line is the Leggett (1992) main-sequence. The dashed line is the completeness limit. The arrow corresponds to $A_V = 1$.

Figure 4: Traces of five representative spectra with $H\alpha$. The spectra are corrected for reddening but not for telluric absorption. UScoCTIO 128 has a $H\alpha$ equivalent width of -130.5.

Figure 5: Absolute values of $H\alpha$ equivalent widths for various clusters. All the $H\alpha$ lines are in emission. The symbols are: (●) Upper-Sco, (*) α -Persei, (◇) σ -Orionis. For clarity, only those objects with $H\alpha$ EqW. less than 50 Å are plotted. The dashed line shows the envelope of the $H\alpha$ equivalent widths for field stars (Prosser et al. 1991; Liebert et al. 1992). The cross indicates the size of the error bars. The $H\alpha$ strength in Upper-Sco is intermediate between that of σ -Orionis and α -Persei.

Figure 6: Color-magnitude diagram of objects observed spectroscopically. All objects have been de-reddened. The dots (●) indicate objects for which the $H\alpha$ equivalent width is above the field envelope (see text). The asterisks (*) indicate objects for which the $H\alpha$ equivalent width is below the field. The upper axis shows the spectral types calculated using the color-spectral type calibration from Kirkpatrick & McCarthy (1994). Also shown are the evolutionary models by D'Antona & Mazzitelli (1994), for masses from 0.20 to

$0.02M_{\odot}$, and isochrones from 1 Myr to 10 Myr. Of those objects observed spectroscopically only two (UScoCTIO 28 and UScoCTIO 162) do not have $H\alpha$.




Table 1. Summary of observations

Site	Date	fields/objects	Observations
CTIO Schmidt	April 1-5, 1998	8 fields 80' by 80'	Imaging R,I,Z
CTIO 1.5 m	February 28 to March 2, 1999	9 objects	Imaging J,H
Lick 3 m	June 21-23 and July 19th, 1999	22 objects	Low-res. Optical Spectra

This figure "upper_sco.fig2a.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/0003316v1>

This figure "upper_sco.fig2b.jpeg" is available in "jpeg" format from:

<http://arXiv.org/ps/astro-ph/0003316v1>

Table 2. RIZ candidate members in the Upper-Scorpius OB Association

Name	R.A. (J2000)	Dec. (J2000)	I	R-I
UScoCTIO 1	16:07:11.39	-17:09:12.2	13.48 ± 0.07	1.21 ± 0.09
UScoCTIO 2	15:53:18.82	-23:06:39.5	13.52 ± 0.07	1.73 ± 0.11
UScoCTIO 3	16:03:50.00	-18:05:45.9	13.55 ± 0.07	2.09 ± 0.10
UScoCTIO 4	15:53:57.22	-23:11:45.2	13.56 ± 0.10	2.05 ± 0.13
UScoCTIO 5	15:59:50.39	-19:44:35.8	13.65 ± 0.07	1.77 ± 0.10
UScoCTIO 6	16:00:07.03	-23:40:47.4	13.67 ± 0.06	1.92 ± 0.11
UScoCTIO 7	16:01:28.78	-25:09:06.0	13.68 ± 0.07	1.47 ± 0.10
UScoCTIO 8	16:01:30.30	-24:21:31.6	13.71 ± 0.08	1.63 ± 0.10
UScoCTIO 9	16:04:39.66	-22:47:05.3	13.72 ± 0.08	1.74 ± 0.10
UScoCTIO 10	15:53:13.02	-26:32:57.3	13.76 ± 0.06	1.58 ± 0.09
UScoCTIO 11	16:00:35.87	-20:11:53.6	13.78 ± 0.07	2.03 ± 0.10
UScoCTIO 12	16:02:04.95	-23:31:07.1	13.81 ± 0.09	1.65 ± 0.11
UScoCTIO 13	15:57:29.72	-22:58:43.2	13.81 ± 0.09	1.82 ± 0.12
UScoCTIO 14	15:39:56.45	-18:29:10.4	13.82 ± 0.04	1.76 ± 0.06
UScoCTIO 15	16:02:23.50	-22:59:32.4	13.87 ± 0.07	2.14 ± 0.10
UScoCTIO 16	16:02:56.42	-22:56:24.1	13.90 ± 0.07	1.88 ± 0.12
UScoCTIO 17	16:06:34.42	-22:55:03.2	13.92 ± 0.07	1.80 ± 0.11
UScoCTIO 18	15:44:05.08	-17:49:47.6	13.93 ± 0.05	1.62 ± 0.07
UScoCTIO 19	16:01:13.60	-25:16:26.8	13.95 ± 0.07	1.81 ± 0.11
UScoCTIO 20	16:01:26.61	-23:01:33.0	13.98 ± 0.07	1.92 ± 0.10
UScoCTIO 21	15:58:36.18	-19:46:12.6	14.00 ± 0.07	1.81 ± 0.11
UScoCTIO 22	15:53:06.47	-26:06:31.4	14.00 ± 0.07	1.65 ± 0.08
UScoCTIO 23	16:03:22.54	-22:38:21.2	14.02 ± 0.07	1.89 ± 0.11
UScoCTIO 24	16:01:35.26	-24:43:35.7	14.04 ± 0.07	1.71 ± 0.11
UScoCTIO 25	15:59:44.32	-20:29:22.6	14.04 ± 0.07	1.85 ± 0.11

Table 2—Continued

Name	R.A. (J2000)	Dec. (J2000)	I	R-I
UScoCTIO 26	16:01:57.66	-22:50:22.0	14.09 ± 0.04	1.38 ± 0.12
UScoCTIO 27	16:01:22.66	-24:07:57.1	14.13 ± 0.07	1.50 ± 0.11
UScoCTIO 28	15:57:53.91	-23:17:52.3	14.14 ± 0.05	1.85 ± 0.10
UScoCTIO 29	16:00:32.38	-25:00:23.1	14.18 ± 0.07	1.73 ± 0.11
UScoCTIO 30	15:58:20.94	-20:05:13.1	14.21 ± 0.08	2.03 ± 0.11
UScoCTIO 31	16:03:23.52	-24:13:19.5	14.22 ± 0.07	1.56 ± 0.11
UScoCTIO 32	16:01:02.53	-20:36:12.5	14.23 ± 0.07	1.50 ± 0.11
UScoCTIO 33	15:58:29.75	-23:10:07.4	14.23 ± 0.07	1.55 ± 0.12
UScoCTIO 34	15:52:35.49	-22:39:06.8	14.23 ± 0.05	1.97 ± 0.13
UScoCTIO 35	16:02:23.34	-22:59:33.2	14.26 ± 0.08	2.05 ± 0.12
UScoCTIO 36	16:01:01.41	-20:58:02.2	14.26 ± 0.07	1.48 ± 0.11
UScoCTIO 37	16:02:54.24	-18:05:29.1	14.26 ± 0.07	2.09 ± 0.11
UScoCTIO 38	16:06:39:02	-22:48:34.1	14.28 ± 0.08	1.81 ± 0.13
UScoCTIO 39	15:58:15.77	-20:21:36.1	14.28 ± 0.07	1.98 ± 0.11
UScoCTIO 40	16:01:23.70	-24:19:47.0	14.30 ± 0.07	2.15 ± 0.11
UScoCTIO 41	16:06:07.74	-18:03:00.8	14.31 ± 0.08	2.07 ± 0.11
UScoCTIO 42	15:52:55.34	-26:04:42.4	14.31 ± 0.06	1.66 ± 0.07
UScoCTIO 43	16:07:50.37	-22:21:01.2	14.34 ± 0.08	1.95 ± 0.10
UScoCTIO 44	16:00:48.15	-23:21:23.9	14.34 ± 0.05	1.76 ± 0.11
UScoCTIO 45	16:02:45.70	-23:04:50.5	14.35 ± 0.07	2.19 ± 0.11
UScoCTIO 46	15:54:02.40	-22:54:57.5	14.36 ± 0.09	1.81 ± 0.10
UScoCTIO 47	16:02:25.86	-22:24:55.7	14.39 ± 0.07	2.06 ± 0.11
UScoCTIO 48	16:04:10.06	-22:34:46.0	14.40 ± 0.07	1.90 ± 0.13
UScoCTIO 49	16:06:08.73	-18:03:14.9	14.40 ± 0.07	2.13 ± 0.10
UScoCTIO 50	16:07:45.10	-22:22:57.0	14.49 ± 0.07	1.80 ± 0.10

Table 2—Continued

Name	R.A. (J2000)	Dec. (J2000)	I	R-I
UScoCTIO 51	15:42:11.19	-18:29:10.4	14.52 ± 0.06	1.87 ± 0.07
UScoCTIO 52	16:00:43.06	-24:30:48.7	14.52 ± 0.07	1.93 ± 0.11
UScoCTIO 53	16:00:26.31	-22:59:40.4	14.54 ± 0.05	2.22 ± 0.12
UScoCTIO 54	16:02:12.71	-23:21:20.5	14.56 ± 0.06	1.94 ± 0.09
UScoCTIO 55	16:02:45.60	-23:04:49.8	14.58 ± 0.60	2.12 ± 0.09
UScoCTIO 56	16:01:40.95	-20:22:08.1	14.62 ± 0.07	2.03 ± 0.11
UScoCTIO 57	16:00:52.03	-23:51:21.8	14.65 ± 0.04	1.74 ± 0.10
UScoCTIO 58	15:58:39.87	-23:07:10.2	14.67 ± 0.08	1.98 ± 0.10
UScoCTIO 59	16:03:01.70	-22:07:52.6	14.70 ± 0.08	2.02 ± 0.12
UScoCTIO 60	16:02:09.50	-23:02:27.0	14.71 ± 0.06	2.08 ± 0.13
UScoCTIO 61	15:41:39.70	-18:37:14.0	14.73 ± 0.07	1.56 ± 0.09
UScoCTIO 62	15:59:17.86	-23:43:05.3	14.76 ± 0.06	2.07 ± 0.07
UScoCTIO 63	15:54:32.30	-26:29:31.2	14.77 ± 0.08	1.71 ± 0.09
UScoCTIO 64	15:59:35.91	-23:23:41.1	14.83 ± 0.05	1.62 ± 0.13
UScoCTIO 65	15:52:39.80	-25:42:26.3	14.85 ± 0.09	1.66 ± 0.10
UScoCTIO 66	16:01:49.66	-23:51:07.4	14.85 ± 0.08	2.18 ± 0.12
UScoCTIO 67	15:59:25.94	-23:05:08.1	14.87 ± 0.09	2.12 ± 0.09
UScoCTIO 68	15:55:30.70	-23:45:19.9	14.90 ± 0.10	1.88 ± 0.12
UScoCTIO 69	16:03:53.90	-25:09:38.4	15.00 ± 0.08	1.97 ± 0.11
UScoCTIO 70	16:02:10.91	-20:07:49.3	15.01 ± 0.07	1.80 ± 0.11
UScoCTIO 71	15:54:42.61	-26:26:25.4	15.05 ± 0.09	1.71 ± 0.11
UScoCTIO 72	16:05:40.89	-17:59:20.7	15.06 ± 0.06	2.25 ± 0.11
UScoCTIO 73	16:00:29.58	-20:22:53.3	15.06 ± 0.07	2.00 ± 0.11
UScoCTIO 74	16:02:14.78	-24:38:31.5	15.06 ± 0.08	2.14 ± 0.11
UScoCTIO 75	16:00:30.17	-23:34:44.7	15.08 ± 0.05	2.17 ± 0.08

Table 2—Continued

Name	R.A. (J2000)	Dec. (J2000)	I	R-I
UScoCTIO 76	16:06:15.94	-22:18:27.3	15.08 ± 0.08	1.87 ± 0.12
UScoCTIO 77	16:02:57.42	-24:42:40.6	15.08 ± 0.08	2.02 ± 0.11
UScoCTIO 78	15:52:27.70	-23:02:36.0	15.09 ± 0.10	1.74 ± 0.14
UScoCTIO 79	15:59:49.74	-23:01:57.8	15.09 ± 0.09	1.88 ± 0.12
UScoCTIO 80	15:58:35.87	-23:48:12.2	15.12 ± 0.07	2.12 ± 0.09
UScoCTIO 81	16:01:54.91	-23:36:07.3	15.13 ± 0.08	2.00 ± 0.11
UScoCTIO 82	16:01:06.82	-20:59:19.0	15.17 ± 0.08	1.73 ± 0.11
UScoCTIO 83	16:04:42.54	-22:54:52.7	15.21 ± 0.08	1.72 ± 0.10
UScoCTIO 84	16:06:26.34	-23:06:10.6	15.22 ± 0.07	1.81 ± 0.10
UScoCTIO 85	15:54:03.51	-23:12:31.0	15.23 ± 0.07	2.23 ± 0.10
UScoCTIO 86	16:02:41.34	-22:48:42.1	15.27 ± 0.08	2.17 ± 0.11
UScoCTIO 87	16:03:38.26	-17:09:19.7	15.29 ± 0.08	2.02 ± 0.10
UScoCTIO 88	15:52:29.26	-23:52:09.2	15.34 ± 0.10	1.89 ± 0.14
UScoCTIO 89	15:59:59.10	-24:38:14.2	15.37 ± 0.08	1.89 ± 0.12
UScoCTIO 90	15:59:30.05	-22:51:25.2	15.43 ± 0.05	1.88 ± 0.10
UScoCTIO 91	15:59:57.94	-24:38:13.3	15.44 ± 0.08	1.91 ± 0.12
UScoCTIO 92	15:56:00.62	-23:28:10.3	15.45 ± 0.11	1.75 ± 0.13
UScoCTIO 93	16:04:14.66	-24:24:42.6	15.46 ± 0.08	2.03 ± 0.12
UScoCTIO 94	16:03:34.81	-24:20:24.3	15.48 ± 0.08	2.12 ± 0.12
UScoCTIO 95	16:04:41.79	-17:41:19.2	15.50 ± 0.08	1.90 ± 0.11
UScoCTIO 96	15:52:48.55	-26:21:43.8	15.52 ± 0.09	1.81 ± 0.10
UScoCTIO 97	15:59:31.39	-23:31:56.4	15.58 ± 0.06	2.00 ± 0.10
UScoCTIO 98	16:02:06.78	-25:12:37.7	15.58 ± 0.08	2.01 ± 0.12
UScoCTIO 99	16:01:35.42	-23:15:55.4	15.59 ± 0.08	2.04 ± 0.11
UScoCTIO 100	16:02:04.13	-20:50:41.5	15.62 ± 0.08	2.33 ± 0.12

Table 2—Continued

Name	R.A. (J2000)	Dec. (J2000)	I	R-I
UScoCTIO 101	15:57:51.92	-20:12:32.3	15.64 ± 0.08	2.05 ± 0.12
UScoCTIO 102	16:02:41.38	-22:48:42.0	15.65 ± 0.10	2.05 ± 0.15
UScoCTIO 103	16:00:44.53	-24:50:33.8	15.67 ± 0.08	2.07 ± 0.12
UScoCTIO 104	15:57:12.66	-23:43:45.3	15.68 ± 0.05	2.08 ± 0.10
UScoCTIO 105	16:05:29.70	-18:07:10.0	15.71 ± 0.08	2.24 ± 0.11
UScoCTIO 106	15:58:26.22	-24:01:12.5	15.74 ± 0.06	2.10 ± 0.07
UScoCTIO 107	15:57:55.97	-20:35:10.6	15.79 ± 0.08	2.20 ± 0.12
UScoCTIO 108	16:05:53.94	-18:18:42.7	15.88 ± 0.07	2.19 ± 0.12
UScoCTIO 109	16:01:19.10	-23:06:38.6	16.06 ± 0.09	2.15 ± 0.12
UScoCTIO 110	16:04:42.82	-23:18:26.3	16.07 ± 0.06	2.30 ± 0.09
UScoCTIO 111	15:59:57.12	-24:11:36.5	16.14 ± 0.08	1.98 ± 0.12
UScoCTIO 112	16:00:26.57	-20:56:32.0	16.14 ± 0.08	2.08 ± 0.12
UScoCTIO 113	15:56:01.10	-23:38:06.3	16.18 ± 0.07	1.94 ± 0.10
UScoCTIO 114	15:52:31.88	-23:53:53.1	16.19 ± 0.06	2.10 ± 0.11
UScoCTIO 115	16:05:16.10	-23:12:23.0	16.24 ± 0.08	1.89 ± 0.10
UScoCTIO 116	15:52:22.65	-23:13:36.2	16.33 ± 0.10	2.11 ± 0.12
UScoCTIO 117	16:04:55.82	-23:07:42.4	16.38 ± 0.08	2.27 ± 0.10
UScoCTIO 118	16:00:13.94	-23:27:11.5	16.38 ± 0.05	1.91 ± 0.08
UScoCTIO 119	16:00:17.99	-22:56:54.5	16.39 ± 0.08	2.15 ± 0.10
UScoCTIO 120	15:59:17.71	-23:14:47.2	16.41 ± 0.07	2.06 ± 0.08
UScoCTIO 121	15:51:47.34	-26:23:47.3	16.46 ± 0.06	2.25 ± 0.07
UScoCTIO 122	16:02:02.29	-20:32:34.7	16.47 ± 0.08	1.91 ± 0.12
UScoCTIO 123	16:04:51.94	-22:24:11.8	16.76 ± 0.08	2.33 ± 0.11
UScoCTIO 124	16:03:09.87	-23:28:46.1	16.76 ± 0.04	2.01 ± 0.10
UScoCTIO 125	16:02:57.86	-24:23:37.4	16.82 ± 0.08	2.23 ± 0.13

Table 2—Continued

Name	R.A. (J2000)	Dec. (J2000)	I	R-I
UScoCTIO 126	16:06:05.78	-22:39:34.8	16.87 ± 0.07	2.09 ± 0.11
UScoCTIO 127	16:04:40.28	-22:54:31.8	16.95 ± 0.07	2.23 ± 0.10
UScoCTIO 128	15:59:11.20	-23:37:59.0	17.09 ± 0.06	2.29 ± 0.07
UScoCTIO 129	15:58:56.47	-23:03:34.4	17.32 ± 0.07	2.08 ± 0.11
UScoCTIO 130	15:59:43.56	-20:14:38.1	17.45 ± 0.09	2.46 ± 0.13
UScoCTIO 131	16:00:19.48	-22:56:28.9	17.48 ± 0.05	2.53 ± 0.12
UScoCTIO 132	15:59:37.74	-22:54:09.5	17.63 ± 0.07	2.34 ± 0.11
UScoCTIO 133	15:54:30.74	-25:36:01.8	17.64 ± 0.08	2.24 ± 0.11
UScoCTIO 134	16:06:47.83	-22:30:36.8	17.76 ± 0.09	2.54 ± 0.11
UScoCTIO 135	15:57:52.33	-19:56:30.6	17.88 ± 0.09	2.71 ± 0.14
UScoCTIO 136	16:05:01.96	-23:21:34.4	18.09 ± 0.08	2.59 ± 0.11
UScoCTIO 137	15:56:47.87	-23:47:44.0	18.21 ± 0.07	2.58 ± 0.24
UScoCTIO 138	16:00:52.19	-20:02:14.0	18.73 ± 0.09	2.53 ± 0.14




Table 3. Spectroscopic observations

Name	H α EqW ^a	SpT ^b	A_V ^c	R-I ^d	J ^f	H ^g	SpT ^h
UScoCTIO 5	-2.7	M4	0.5	1.63 ± 0.10	-	-	-
UScoCTIO 18	< -1.0	M4	0.0	1.62 ± 0.07	-	-	-
UScoCTIO 40	-8.0	M5	0.5	2.01 ± 0.11	-	-	-
UScoCTIO 53	-11.0	M5	1.0	1.95 ± 0.12	-	-	-
UScoCTIO 55	-9.5	M5.5	0.5	1.98 ± 0.09	-	-	-
UScoCTIO 56	-9.1	M5	0.5	1.89 ± 0.11	-	-	-
UScoCTIO 60	-6.2	M4	1.5	1.67 ± 0.13	-	-	-
UScoCTIO 63	-5.7	M4	0.0	1.71 ± 0.09	-	-	-
UScoCTIO 66	-7.4	M6	0.0	2.18 ± 0.12	-	-	-
UScoCTIO 67	-16.0	M5.5	0.5	1.98 ± 0.09	-	-	-
UScoCTIO 75	-12.6	M6	0.0	2.17 ± 0.08	-	-	-
UScoCTIO 80	-8.7	M4	2.0	1.58 ± 0.09	-	-	-
UScoCTIO 85	-6.0	M6	0.0	2.23 ± 0.10	13.025 ± 0.03	12.371 ± 0.05	M5-6
UScoCTIO 99	-9.5	M5	0.5	1.90 ± 0.11	-	-	-
UScoCTIO 100	-16.1	M7	0.0	2.33 ± 0.12	-	-	-
UScoCTIO 101	-6.9	M5	0.0	2.05 ± 0.12	-	-	-
UScoCTIO 104	-10.2	M5	0.0	2.08 ± 0.10	13.529 ± 0.06	12.975 ± 0.09	M5-6
UScoCTIO 109	-19.0	M6	0.0	2.15 ± 0.12	-	-	-
UScoCTIO 112	-12.2	M5.5	0.0	2.08 ± 0.12	-	-	-
UScoCTIO 114	-	-	-	2.10 ± 0.11	14.321 ± 0.01	13.538 ± 0.01	M5-6
UScoCTIO 121	-10.0	M6	0.0	2.25 ± 0.07	14.404 ± 0.01	13.798 ± 0.01	M5-6
UScoCTIO 128	-130.5	M7	0.0	2.29 ± 0.07	14.413 ± 0.06	13.780 ± 0.07	M7-8
UScoCTIO 130	-	-	-	2.46 ± 0.13	15.740 ± 0.03	14.979 ± 0.03	M7-8
UScoCTIO 131	-	-	-	2.53 ± 0.12	15.740 ± 0.03	14.979 ± 0.03	M6.5
UScoCTIO 132	< -1.0	M7	0.0	2.34 ± 0.11	14.355 ± 0.06	13.589 ± 0.07	M8-9

Table 3—Continued

Name	H α EqW ^a	SpT ^b	A_V ^c	R-I ^d	J ^f	H ^g	SpT ^h
UScoCTIO 137	-	-	-	2.58 ± 0.24	15.740 ± 0.03	14.979 ± 0.03	M7

^aEquivalent width of the H α line in Å. The upper limits are non-detections.

^bSpectral type. The accuracy is half a spectral subclass.

^cReddening. The accuracy is half a magnitude.

^dR-I color, with reddening correction applied (if known) from Cardelli, Clayton & Mathis (1989).

^eSpectral type from the R-I color (Kirkpatrick & McCarthy 1994).

^fCIRIM photometry, J magnitude.

^gCIRIM photometry, H magnitude.

^hSpectral type from the I-J color (Kirkpatrick & McCarthy 1994).

